CONTENTS OF HEREDITAS, VOLUME 136 (2002)

207 Andersson, S. and Waldmann, P.
Inbreeding depression in a rare plant, *Scabiosa canescens* (Dipsacaceae).

116 Assefa, K., Tefera, H. and Merker, A.
Variation and inter-relationships of quantitative traits in tef (*Eragrostis tef* (Zucc.) Trotter) germplasm from western and southern Ethiopia.

51 Bertolotto, C.E.V., Pellegrino, K.C.M., Rodrigues, M.T. and Yonenaga-Yassuda, Y.
Comparative cytogenetics and supernumerary chromosomes in the Brazilian lizard genus *Enyalius* (Squamata, Polychrotidae).

97 Bione, N.C.P., Pagliarini, M.S. and de Almeida, L.A.
A new and distinct male-sterile, female-fertile desynaptic mutant in soybean (*Glycine max*).

254 Bureš, S., Nádvorník, P. and Sætre, G.-P.
Hybridization and apparent hybridization between meadow pipit (*Anthus pratensis*) and water pipit (*A. spinoletta*).

130 Catanesi, C.I., Vidal-Rioja, L., Crisci, J.V. and Zambelli, A.
Phylogenetic relationships among Robertsonian karyomorphs of *Graomys griseflavus* (Rodentia: Muridae) by mitochondrial cytochrome *b* DNA sequencing.

231 Cheng, Z.C., Lu, B.-R., Baldwin, B.S., Sameshima, K. and Chen, J.-K.
Comparative studies of genetic diversity in kenaf (*Hibiscus cannabinus* L.) varieties based on analysis of agronomic and RAPD data.

126 Chung, H.-S., Um, J.-Y., Kim, M.-S., Hong, S.-H., Kim, S.-M., Kim, H.y-K., Park, S.-J., Kim, S.-C., Hwang, W.-J. and Kim, H.-M.
Determination of the site of origin of *Pinella ternata* roots based on RAPD analysis and PCR-RFLP.

159 De Carvalho, R. and Guerra, M.
Cytogenetics of *Manihot esculenta* Crantz (cassava) and eight related species.

59 Ellneskog-Staam, P. and Merker, A.
Chromosome composition, stability and fertility of alloploids between *Triticum turgidum* var. *carthlicum* and *Thinopyrum junceiforme*.

151 Hanada, H.
G and C banding show structural differences between the Z and W chromosomes in the frog *Buergeria buergeri*.

212 Huang, X.Q., Hsam, S.L.K. and Zeller, F.J.
Chromosomal location of genes for resistance to powdery mildew in Chinese wheat lines Jieyan 94-1-1 and Siyan 94-1-2.

184 Jensen, L.H., Enghoff, H., Frydenberg, J. and Parker, Jr, E.D.
Genetic diversity and the phylogeography of parthenogenesis: comparing bisexual and thelytokous populations of *Nemasoma varicorne* (Diplopoda: Nemasomatidae) in Denmark.

169 Kankare, M., Várkonyi, G. and Saccheri, I.
Genetic differentiation between alternate-year cohorts of *Xestia tecta* (Lepidoptera, Noctuidae) in Finnish Lapland.

219 Kärkkäinen, K. and Ågren, J.
Genetic basis of trichome production in *Arabidopsis lyrata*.

195 Kawahara, T., Taketa, S. and Murai, K.
Differential effects of cultivated and wild barley 5H chromosomes on heading characters in wheat-barley chromosome addition lines.

247 Keyghobadi, N., Roland, J. and Strobeck, C.
Isolation of novel microsatellite loci in the Rocky Mountain apollo butterfly, *Parnassius smintheus*.

144 Kulak, S., Hasterok, R. and Maluszynska, J.
Karyotyping of *Brassica* amphidiploids using 5S and 25S rDNA as chromosome markers.

108 Liljeroth, E. and Bryngelsson, T.
Earlier onset of DNA fragmentation in leaves of wheat compared to barley and rye.

45 Lind-Halldén, C., Halldén, C. and Säll, T.
Genetic variation in *Arabidopsis suecica* and its parental species *A. arenosa* and *A. thaliana*.

67 Liu, F., Sun, G.-L., Salomon, B. and Von Bothmer, R.
Characterization of genetic diversity in core collection accessions of wild barley, *Hordeum vulgare* ssp. *spontaneum*.

104 Lomholt, B., Christensen, K. and Frederiksen, S.
Guinea pig (*Cavio cambayo*) 5S rRNA genes map to 7q2, 20q2 and 30q2 shown by an R-banded karyotype with PNA-FISH.

251 Luque, C., Legal, L., Staudter, H., Gers, C. and Wink, M.
ISSR (Inter Simple Sequence Repeats) as genetic markers in Noctuids (Lepidoptera).

7 Nokkala, S., Laukkanen, A. and Nokkala, C.
Mitotic and meiotic chromosomes in *Somatochlora metallica* (Cordulidae, Odonata). The absence of localized centromeres and inverted meiosis.

240 O'Grady, P.M., Durando, C.M., Heed, W.B., Wasserman, M., Etges, W. and DeSalle, R.
Genetic divergence within the *Drosophila mayaguana* subcluster, a closely related triad of Caribbean species in the *repleta* species group.

75 Pecsenye, K. and Saura, A.
Structure of variation in enzyme activity in natural *Drosophila melanogaster* populations.

29 Persson, K. and Bothmer, von R.
Genetic diversity amongst landraces of rye (*Secale cereale* L.) from northern Europe.

201 Peusha, H., Lebedeva, T., Priilinn, O. and Enno, T.
Genetic analysis of durable powdery mildew resistance in a common wheat line.

89 Portin, P.
General outlines of the molecular genetics of the Notch signalling pathway in *Drosophila melanogaster*: a review.

227 Porto-Foresti, F., Oliveira, C., Tabata, Y.A., Rigolino, M.G. and Foresti, F.
NORs inheritance analysis in crossings including individuals from two stocks of rainbow trout (*Oncorhynchus mykiss*).

13 Rank, J., Lopez, L.C., Nielsen, M.H. and Moretton, J.
Genotoxicity of maleic hydrazide, acridine and DEHP in Allium cepa root cells performed by two different laboratories.

1 Rasmuson, M.
The genotype-phenotype link.

177 Rasmuson, M.
Fluctuating asymmetry — indicator of what?

155 Remis, M.I.
Differential adult survival and phenotypic variation in a chromosomally polymorphic population of the grasshopper *Sinipta dalmani*.

19 Rocha, M.P., Pompolo, S.D.G., Derdam, J.A., Fernandes, A. and Campos, L.A.O.
DNA characterization and karyotypic evolution in the bee genus *Melipona* (Hymenoptera, Meliponini).

137 Santos, N., Fagundes, V., Yoneyaga-Yassuda, Y. and Souza, M.J.
Localization of rRNA genes in Phyllostomidae bats reveals silent NORs in *Artibeus cinerus*.

39 Yang, H., Ren, X., Weng, Q., Zhu, L. and He, G.
Molecular mapping and genetic analysis of a rice brown planthopper (*Nilaparvata lugens* Stål) resistance gene.

85 Zhu, B., Feng, Z., Qu, A., Gao, H., Zhang, Y., Sun, D., Song, W. and Saura, A.
The karyotype of the caudate amphibian *Andrias davidianus*.